



# Normal Criteria for Families of Meromorphic Functions

Gerd Dethloff, Tran van Tan, Nguyen van Thin

## ► To cite this version:

Gerd Dethloff, Tran van Tan, Nguyen van Thin. Normal Criteria for Families of Meromorphic Functions. Journal of Mathematical Analysis and Applications, 2012, 411, pp.675-683. hal-00753121v2

**HAL Id: hal-00753121**

**<https://hal.science/hal-00753121v2>**

Submitted on 8 Dec 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# NORMAL CRITERIA FOR FAMILIES OF MEROMORPHIC FUNCTIONS

GERD DETHLOFF<sup>a</sup> AND TRAN VAN TAN<sup>b, 1</sup> AND NGUYEN VAN THIN<sup>c</sup>

<sup>a</sup> Université de Brest, LMBA, UMR CNRS 6205,  
6, avenue Le Gorgeu - C.S. 93837, 29238 Brest Cedex 3, France

<sup>b</sup> Department of Mathematics, Hanoi National University of Education,  
136 Xuan Thuy Street, Cau Giay, Hanoi, Vietnam

<sup>c</sup> Department of Mathematics, Thai Nguyen University of Education,  
Luong Ngoc Quyen Street, Thai Nguyen City, Vietnam

## Abstract

By using Nevanlinna theory, we prove some normality criteria for a family of meromorphic functions under a condition on differential polynomials generated by the members of the family.

*Keywords:* Meromorphic function, Normal family, Nevanlinna theory.

Mathematics Subject Classification 2010: 30D35.

## 1 Introduction

Let  $D$  be a domain in the complex plane  $\mathbb{C}$  and  $\mathcal{F}$  be a family of meromorphic functions in  $D$ . The family  $\mathcal{F}$  is said to be normal in  $D$ , in the sense of Montel, if for any sequence  $\{f_v\} \subset \mathcal{F}$ , there exists a subsequence  $\{f_{v_i}\}$  such that  $\{f_{v_i}\}$  converges spherically locally uniformly in  $D$ , to a meromorphic function or  $\infty$ .

In 1989, Schwick proved:

---

<sup>1</sup>Corresponding author.

E-mail addresses: <sup>a</sup>gerd.dethloff@univ-brest.fr, <sup>b</sup>tranvantanh@yahoo.com,  
<sup>c</sup>thinmath@gmail.com

The second named author is currently Regular Associate Member of ICTP, Trieste, Italy. This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED).

**Theorem A** ([6], Theorem 3.1). *Let  $k, n$  be positive integers such that  $n \geq k + 3$ . Let  $\mathcal{F}$  be a family of meromorphic functions in a complex domain  $D$  such that for every  $f \in \mathcal{F}$ ,  $(f^n)^{(k)}(z) \neq 1$  for all  $z \in D$ . Then  $\mathcal{F}$  is normal on  $D$ .*

**Theorem B** ([6], Theorem 3.2). *Let  $k, n$  be positive integers such that  $n \geq k + 1$ . Let  $\mathcal{F}$  be a family of entire functions in a complex domain  $D$  such that for every  $f \in \mathcal{F}$ ,  $(f^n)^{(k)}(z) \neq 1$  for all  $z \in D$ . Then  $\mathcal{F}$  is normal on  $D$ .*

The following normality criterion was established by Pang and Zalcman [7] in 1999:

**Theorem C** ([7]). *Let  $n$  and  $k$  be natural numbers and  $\mathcal{F}$  be a family of holomorphic functions in a domain  $D$  all of whose zeros have multiplicity at least  $k$ . Assume that  $f^n f^{(k)} - 1$  is non-vanishing for each  $f \in \mathcal{F}$ . Then  $\mathcal{F}$  is normal in  $D$ .*

The main purpose of this paper is to establish some normality criteria for the case of more general differential polynomials. Our main results are as follows:

**Theorem 1.** *Take  $q$  ( $q \geq 1$ ) distinct nonzero complex values  $a_1, \dots, a_q$ , and  $q$  positive integers (or  $+\infty$ )  $\ell_1, \dots, \ell_q$ . Let  $n$  be a nonnegative integer, and let  $n_1, \dots, n_k, t_1, \dots, t_k$  be positive integers ( $k \geq 1$ ). Let  $\mathcal{F}$  be a family of meromorphic functions in a complex domain  $D$  such that for every  $f \in \mathcal{F}$  and for every  $m \in \{1, \dots, q\}$ , all zeros of  $f^n (f^{n_1})^{(t_1)} \dots (f^{n_k})^{(t_k)} - a_m$  have multiplicity at least  $\ell_m$ . Assume that*

a)  $n_j \geq t_j$  for all  $1 \leq j \leq k$ , and  $\ell_i \geq 2$  for all  $1 \leq i \leq q$ ,

b)  $\sum_{i=1}^q \frac{1}{\ell_i} < \frac{qn-2+\sum_{j=1}^k q(n_j-t_j)}{n+\sum_{j=1}^k (n_j+t_j)}$ .

*Then  $\mathcal{F}$  is a normal family.*

Take  $q = 1$  and  $\ell_1 = +\infty$ , we get the following corollary of Theorem 1:

**Corollary 2.** *Let  $a$  be a nonzero complex value, let  $n$  be a nonnegative integer, and  $n_1, \dots, n_k, t_1, \dots, t_k$  be positive integers. Let  $\mathcal{F}$  be a family of meromorphic functions in a complex domain  $D$  such that for every  $f \in \mathcal{F}$ ,  $f^n (f^{n_1})^{(t_1)} \dots (f^{n_k})^{(t_k)} - a$  is nowhere vanishing on  $D$ . Assume that*

a)  $n_j \geq t_j$  for all  $1 \leq j \leq k$ ,

b)  $n + \sum_{j=1}^k n_j \geq 3 + \sum_{j=1}^k t_j$ .

*Then  $\mathcal{F}$  is normal on  $D$ .*

We remark that in the case where  $n \geq 3$ , condition a) in the above corollary implies condition b); and in the case where  $n = 0$  and  $k = 1$ , Corollary 2 gives Theorem A.

For the case of entire functions, we shall prove the following result:

**Theorem 3.** Take  $q$  ( $q \geq 1$ ) distinct nonzero complex values  $a_1, \dots, a_q$ , and  $q$  positive integers (or  $+\infty$ )  $\ell_1, \dots, \ell_q$ . Let  $n$  be a nonnegative integer, and let  $n_1, \dots, n_k, t_1, \dots, t_k$  be positive integers ( $k \geq 1$ ). Let  $\mathcal{F}$  be a family of holomorphic functions in a complex domain  $D$  such that for every  $f \in \mathcal{F}$  and for every  $m \in \{1, \dots, q\}$ , all zeros of  $f^n(f^{n_1})^{(t_1)} \dots (f^{n_k})^{(t_k)} - a_m$  have multiplicity at least  $\ell_m$ . Assume that

a)  $n_j \geq t_j$  for all  $1 \leq j \leq k$ , and  $\ell_i \geq 2$  for all  $1 \leq i \leq q$ ,

b)  $\sum_{i=1}^q \frac{1}{\ell_i} < \frac{qn-1+\sum_{j=1}^k q(n_j-t_j)}{n+\sum_{j=1}^k n_j}$ .

Then  $\mathcal{F}$  is a normal family.

Take  $q = 1$  and  $\ell_1 = +\infty$ , Theorem 3 gives the following generalization of Theorem B, except for the case  $n = k + 1$ . So for the latter case, we add a new proof of Theorem B in the Appendix which is slightly simpler than the original one.

**Corollary 4.** Let  $a$  be a nonzero complex value, let  $n$  be a nonnegative integer, and  $n_1, \dots, n_k, t_1, \dots, t_k$  be positive integers. Let  $\mathcal{F}$  be a family of holomorphic functions in a complex domain  $D$  such that for every  $f \in \mathcal{F}$ ,  $f^n(f^{n_1})^{(t_1)} \dots (f^{n_k})^{(t_k)} - a$  is nowhere vanishing on  $D$ . Assume that

a)  $n_j \geq t_j$  for all  $1 \leq j \leq k$ ,

b)  $n + \sum_{j=1}^k n_j \geq 2 + \sum_{j=1}^k t_j$ .

Then  $\mathcal{F}$  is normal on  $D$ .

In the case where  $n \geq 2$ , condition a) in the above corollary implies condition b).

**Remark 5.** Our above results remain valid if the monomial  $f^n(f^{n_1})^{(t_1)} \dots (f^{n_k})^{(t_k)}$  is replaced by the following polynomial

$$f^n(f^{n_1})^{(t_1)} \dots (f^{n_k})^{(t_k)} + \sum_I c_I f^{n_I} (f^{n_{1I}})^{(t_{1I})} \dots (f^{n_{kI}})^{(t_{kI})},$$

where  $c_I$  is a holomorphic function on  $D$ , and  $n_I, n_{jI}, t_{jI}$  are nonnegative integers satisfying

$$\alpha_I := \frac{\sum_{j=1}^k t_{jI}}{n_I + \sum_{j=1}^k n_{jI}} < \alpha := \frac{\sum_{j=1}^k t_j}{n + \sum_{j=1}^k n_j}.$$

## 2 Some notations and results of Nevanlinna theory

Let  $\nu$  be a divisor on  $\mathbb{C}$ . The counting function of  $\nu$  is defined by

$$N(r, \nu) = \int_1^r \frac{n(t)}{t} dt \quad (r > 1), \text{ where } n(t) = \sum_{|z| \leq t} \nu(z).$$

For a meromorphic function  $f$  on  $\mathbb{C}$  with  $f \not\equiv \infty$ , denote by  $\nu_f$  the pole divisor of  $f$ , and the divisor  $\bar{\nu}_f$  is defined by  $\bar{\nu}_f(z) := \min\{\nu_f(z), 1\}$ . Set  $N(r, f) := N(r, \nu_f)$  and  $\bar{N}(r, f) := N(r, \bar{\nu}_f)$ .

The proximity function of  $f$  is defined by

$$m(r, f) = \frac{1}{2\pi} \int_0^{2\pi} \log^+ |f(re^{i\theta})| d\theta,$$

where  $\log^+ x = \max\{\log x, 0\}$  for  $x \geq 0$ .

The characteristic function of  $f$  is defined by

$$T(r, f) := m(r, f) + N(r, f).$$

We state the Lemma on Logarithmic Derivative, the First and Second Main Theorems of Nevanlinna theory.

**LEMMA ON LOGARITHMIC DERIVATIVE.** *Let  $f$  be a nonconstant meromorphic function on  $\mathbb{C}$ , and let  $k$  be a positive integer. Then the equality*

$$m(r, \frac{f^{(k)}}{f}) = o(T(r, f))$$

*holds for all  $r \in [1, \infty)$  excluding a set of finite Lebesgue measure.*

**FIRST MAIN THEOREM.** *Let  $f$  be a meromorphic functions on  $\mathbb{C}$  and  $a$  be a complex number. Then*

$$T(r, \frac{1}{f-a}) = T(r, f) + O(1).$$

**SECOND MAIN THEOREM.** *Let  $f$  be a nonconstant meromorphic function on  $\mathbb{C}$ . Let  $a_1, \dots, a_q$  be  $q$  distinct values in  $\mathbb{C}$ . Then*

$$(q-1)T(r, f) \leq \overline{N}(r, f) + \sum_{i=1}^q \overline{N}\left(r, \frac{1}{f-a_i}\right) + o(T(r, f)),$$

for all  $r \in [1, \infty)$  excluding a set of finite Lebesgue measure.

### 3 Proof of our results

To prove our results, we need the following lemmas:

**Lemma 6** (Zalcman's Lemma, see [8]). *Let  $\mathcal{F}$  be a family of meromorphic functions defined in the unit disc  $\Delta$ . Then if  $\mathcal{F}$  is not normal at a point  $z_0 \in \Delta$ , there exist, for each real number  $\alpha$  satisfying  $-1 < \alpha < 1$ ,*

- 1) *a real number  $r$ ,  $0 < r < 1$ ,*
- 2) *points  $z_n$ ,  $|z_n| < r$ ,  $z_n \rightarrow z_0$ ,*
- 3) *positive numbers  $\rho_n$ ,  $\rho_n \rightarrow 0^+$ ,*
- 4) *functions  $f_n$ ,  $f_n \in \mathcal{F}$*

*such that*

$$g_n(\xi) = \frac{f_n(z_n + \rho_n \xi)}{\rho_n^\alpha} \rightarrow g(\xi)$$

*spherically uniformly on compact subsets of  $\mathbb{C}$ , where  $g(\xi)$  is a non-constant meromorphic function and  $g^\#(\xi) \leq g^\#(0) = 1$ . Moreover, the order of  $g$  is not greater than 2. Here, as usual,  $g^\#(z) = \frac{|g'(z)|}{1+|g(z)|^2}$  is the spherical derivative.*

**Lemma 7** (see [2]). *Let  $g$  be an entire function and  $M$  is a positive constant. If  $g^\#(\xi) \leq M$  for all  $\xi \in \mathbb{C}$ , then  $g$  has order at most one.*

**Remark 8.** *In Lemma 6, if  $\mathcal{F}$  is a family of holomorphic functions, then by Hurwitz theorem,  $g$  is a holomorphic function. Therefore, by Lemma 7, the order of  $g$  is not greater than 1.*

We consider a nonconstant meromorphic function  $g$  in the complex plane  $\mathbb{C}$ , and its first  $p$  derivatives. A differential polynomial  $P$  of  $g$  is defined by

$$P(z) := \sum_{i=1}^n \alpha_i(z) \prod_{j=0}^p (g^{(j)}(z))^{S_{ij}},$$

where  $S_{ij}$  ( $1 \leq i \leq n$ ,  $0 \leq j \leq p$ ) are nonnegative integers, and  $\alpha_i \neq 0$  ( $1 \leq i \leq n$ ) are small (with respect to  $g$ ) meromorphic functions. Set

$$d(P) := \min_{1 \leq i \leq n} \sum_{j=0}^p S_{ij} \text{ and } \theta(P) := \max_{1 \leq i \leq n} \sum_{j=0}^p j S_{ij}.$$

In 2002, J. Hinchliffe [5] generalized theorems of Hayman [3] and Chuang [1] and obtained the following result:

**Proposition 9.** *Let  $g$  be a transcendental meromorphic function, let  $P(z)$  be a non-constant differential polynomial in  $g$  with  $d(P) \geq 2$ . Then*

$$T(r, g) \leq \frac{\theta(P) + 1}{d(P) - 1} \bar{N}(r, \frac{1}{g}) + \frac{1}{d(P) - 1} \bar{N}(r, \frac{1}{P - 1}) + o(T(r, g)),$$

for all  $r \in [1, +\infty)$  excluding a set of finite Lebesgue measure.

In order to prove our results, we now give the following generalization of the above result:

**Lemma 10.** *Let  $a_1, \dots, a_q$  be distinct nonzero complex numbers. Let  $g$  be a nonconstant meromorphic function, let  $P(z)$  be a nonconstant differential polynomial in  $g$  with  $d(P) \geq 2$ . Then*

$$T(r, g) \leq \frac{q\theta(P) + 1}{qd(P) - 1} \bar{N}(r, \frac{1}{g}) + \frac{1}{qd(P) - 1} \sum_{j=1}^q \bar{N}(r, \frac{1}{P - a_j}) + o(T(r, g)),$$

for all  $r \in [1, +\infty)$  excluding a set of finite Lebesgue measure.

Moreover, in the case where  $g$  is an entire function, we have

$$T(r, g) \leq \frac{q\theta(P) + 1}{qd(P)} \bar{N}(r, \frac{1}{g}) + \frac{1}{qd(P)} \sum_{j=1}^q \bar{N}(r, \frac{1}{P - a_j}) + o(T(r, g)),$$

for all  $r \in [1, +\infty)$  excluding a set of finite Lebesgue measure.

*Proof.* For any  $z$  such that  $|g(z)| \leq 1$ , since  $\sum_{j=0}^p S_{ij} \geq d(P)$  ( $1 \leq i \leq n$ ), we have

$$\begin{aligned} \frac{1}{|g(z)|^{d(P)}} &= \frac{1}{|P(z)|} \cdot \frac{|P(z)|}{|g(z)|^{d(P)}} \\ &\leq \frac{1}{|P(z)|} \cdot \sum_{i=1}^n (|\alpha_i(z)| \prod_{j=0}^p |\frac{g^{(j)}(z)}{g(z)}|^{S_{ij}}). \end{aligned}$$

This implies that for all  $z \in \mathbb{C}$ ,

$$\log^+ \frac{1}{|g(z)|^{d(P)}} \leq \log^+ \left( \frac{1}{|P(z)|} \cdot \sum_{i=1}^n (|\alpha_i(z)| \prod_{j=0}^p \left| \frac{g^{(j)}(z)}{g(z)} \right|^{S_{ij}}) \right).$$

Therefore, by the Lemma on Logarithmic Derivative and by the First Main Theorem, we have

$$\begin{aligned} d(P)m(r, \frac{1}{g}) &\leq m(r, \frac{1}{P}) + o(T(r, g)) = T(r, \frac{1}{P}) - N(r, \frac{1}{P}) + o(T(r, g)) \\ &= T(r, P) - N(r, \frac{1}{P}) + o(T(r, g)). \end{aligned}$$

On the other hand, by the Second Main Theorem (used with the  $q+1$  different values  $0, a_1, \dots, a_q$ ) we have

$$qT(r, P) \leq \bar{N}(r, P) + \bar{N}(r, \frac{1}{P}) + \sum_{j=1}^q \bar{N}(r, \frac{1}{P - a_j}) + o(T(r, g)),$$

Hence,

$$\begin{aligned} d(P)m(r, \frac{1}{g}) &\leq \frac{1}{q} (\bar{N}(r, P) + \bar{N}(r, \frac{1}{P}) + \sum_{j=1}^q \bar{N}(r, \frac{1}{P - a_j})) \\ &\quad - N(r, \frac{1}{P}) + o(T(r, g)). \end{aligned}$$

Therefore, by the First Main Theorem, we have

$$\begin{aligned} d(P)T(r, g) &= d(P)T(r, \frac{1}{g}) + O(1) \\ &= d(P)m(r, \frac{1}{g}) + d(P)N(r, \frac{1}{g}) + O(1) \\ &\leq \frac{1}{q} (\bar{N}(r, P) + \bar{N}(r, \frac{1}{P}) + \sum_{j=1}^q \bar{N}(r, \frac{1}{P - a_j})) \\ &\quad + d(P)N(r, \frac{1}{g}) - N(r, \frac{1}{P}) + o(T(r, g)). \end{aligned} \tag{3.1}$$

We have

$$\frac{1}{g^{d(P)}} = \frac{1}{P(z)} \sum_{i=1}^n (\alpha_i g^{(\sum_{j=0}^p S_{ij}) - d(P)} \prod_{j=0}^p (\frac{g^{(j)}}{g})^{S_{ij}}).$$



(note that  $(\sum_{j=0}^p S_{ij}) - d(P) \geq 0$ ). Therefore,

$$\begin{aligned} d(P)\nu_{\frac{1}{g}} &\leq \nu_{\frac{1}{P}} + \max_{1 \leq i \leq n} \{\nu_{\alpha_i} + \sum_{j=0}^p j S_{ij} \bar{\nu}_{\frac{1}{g}}\} \\ &\leq \nu_{\frac{1}{P}} + \sum_{i=1}^n \nu_{\alpha_i} + \theta(P) \bar{\nu}_{\frac{1}{g}}, \end{aligned}$$

where  $\nu_{\phi}$  is the pole divisor of the meromorphic  $\phi$  and  $\bar{\nu}_{\phi} := \min\{\nu_{\phi}, 1\}$ . This implies,

$$d(P)\nu_{\frac{1}{g}} - \nu_{\frac{1}{P}} + \frac{1}{q} \bar{\nu}_{\frac{1}{P}} \leq (\theta(P) + \frac{1}{q}) \bar{\nu}_{\frac{1}{g}} + \sum_{i=1}^n \nu_{\alpha_i},$$

(note that for any  $z_0$ , if  $\nu_{\frac{1}{g}}(z_0) = 0$  then  $d(P)\nu_{\frac{1}{g}}(z_0) - \nu_{\frac{1}{P}}(z_0) + \frac{1}{q} \bar{\nu}_{\frac{1}{P}}(z_0) \leq 0$ ).

Then,

$$\begin{aligned} d(P)N(r, \frac{1}{g}) - N(r, \frac{1}{P}) + \frac{1}{q} \bar{N}(r, \frac{1}{P}) &\leq (\theta(P) + \frac{1}{q}) \bar{N}(r, \frac{1}{g}) + \sum_{i=1}^n N(r, \alpha_i) \\ &= (\theta(P) + \frac{1}{q}) \bar{N}(r, \frac{1}{g}) + o(T(r, g)). \end{aligned}$$

Combining with (3.1), we have

$$d(P)T(r, g) \leq \frac{1}{q} (\bar{N}(r, P) + \sum_{j=1}^q \bar{N}(r, \frac{1}{P - a_j})) + (\theta(P) + \frac{1}{q}) \bar{N}(r, \frac{1}{g}) + o(T(r, g)).$$

On the other hand, by the definition of the differential polynomial  $P$ ,  $\text{Pole}(P) \subset \cup_{i=1}^n \text{Pole}(\alpha_i) \cup \text{Pole}(g)$ . Hence (since  $\bar{N}(r, \alpha_i) \leq T(r, \alpha_i) = o(T(r, g))$  for  $i = 1, \dots, n$ ), we get

$$\begin{aligned} d(P)T(r, g) &\leq \frac{1}{q} (\bar{N}(r, g) + \sum_{j=1}^q \bar{N}(r, \frac{1}{P - a_j})) + (\theta(P) + \frac{1}{q}) \bar{N}(r, \frac{1}{g}) + o(T(r, g)) \\ &\leq \frac{1}{q} (T(r, g) + \sum_{j=1}^q \bar{N}(r, \frac{1}{P - a_j})) + (\theta(P) + \frac{1}{q}) \bar{N}(r, \frac{1}{g}) + o(T(r, g)). \end{aligned} \tag{3.2}$$

Therefore,

$$T(r, g) \leq \frac{q\theta(P) + 1}{qd(P) - 1} \overline{N}(r, \frac{1}{g}) + \frac{1}{qd(P) - 1} \sum_{j=1}^q \overline{N}(r, \frac{1}{P - a_j}) + o(T(r, g)).$$

In the case where  $g$  is an entire function, the first inequality in (3.2) becomes

$$d(P)T(r, g) \leq \frac{1}{q} \sum_{j=1}^q \overline{N}(r, \frac{1}{P - a_j}) + (\theta(P) + \frac{1}{q}) \overline{N}(r, \frac{1}{g}) + o(T(r, g)).$$

This implies that

$$T(r, g) \leq \frac{\theta(P)q + 1}{qd(P)} \overline{N}(r, \frac{1}{g}) + \frac{1}{qd(P)} \sum_{j=1}^q \overline{N}(r, \frac{1}{P - a_j}) + o(T(r, g)).$$

We have completed the proof of Lemma 10.  $\square$

**Proof of Theorem 1.** Without loss the generality, we may assume that  $D$  is the unit disc. Suppose that  $\mathcal{F}$  is not normal at  $z_0 \in D$ . By Lemma 6, for  $\alpha = \frac{\sum_{j=1}^k t_j}{n + \sum_{j=1}^k n_j}$  there exist

- 1) a real number  $r$ ,  $0 < r < 1$ ,
- 2) points  $z_v$ ,  $|z_v| < r$ ,  $z_v \rightarrow z_0$ ,
- 3) positive numbers  $\rho_v$ ,  $\rho_v \rightarrow 0^+$ ,
- 4) functions  $f_v$ ,  $f_v \in \mathcal{F}$

such that

$$g_v(\xi) = \frac{f_v(z_v + \rho_v \xi)}{\rho_v^\alpha} \rightarrow g(\xi) \quad (3.3)$$

spherically uniformly on compact subsets of  $\mathbb{C}$ , where  $g(\xi)$  is a non-constant meromorphic function and  $g^\#(\xi) \leq g^\#(0) = 1$ .

On the other hand,

$$\begin{aligned} (g_v^{n_j}(\xi))^{(t_j)} &= \left( \left( \frac{f_v(z_v + \rho_v \xi)}{\rho_v^\alpha} \right)^{n_j} \right)^{(t_j)} \\ &= \frac{1}{\rho_v^{n_j \alpha - t_j}} (f_v^{n_j})^{(t_j)}(z_v + \rho_v \xi). \end{aligned}$$

Therefore, by the definition of  $\alpha$  and by (4.1), we have

$$\begin{aligned} f_v^n(z_v + \rho_v \xi) (f_v^{n_1})^{(t_1)}(z_v + \rho_v \xi) \cdots (f_v^{n_k})^{(t_k)}(z_v + \rho_v \xi) \\ = g_v^n(\xi) (g_v^{n_1}(\xi))^{(t_1)} \cdots (g_v^{n_k}(\xi))^{(t_k)} \rightarrow g^n(\xi) (g^{n_1}(\xi))^{(t_1)} \cdots (g^{n_k}(\xi))^{(t_k)} \end{aligned} \quad (3.4)$$

spherically uniformly on compact subsets of  $\mathbb{C}$ .

Now, we prove the following claim:

**Claim:**  $g^n(\xi) (g^{n_1}(\xi))^{(t_1)} \cdots (g^{n_k}(\xi))^{(t_k)}$  is non-constant.

Since  $g$  is non-constant and  $n_j \geq t_j$  ( $j = 1, \dots, k$ ), it easy to see that  $(g^{n_j}(\xi))^{(t_j)} \not\equiv 0$ , for all  $j \in \{1, \dots, k\}$ . Hence,  $g^n(\xi) (g^{n_1}(\xi))^{(t_1)} \cdots (g^{n_k}(\xi))^{(t_k)} \not\equiv 0$ .

Suppose that  $g^n(\xi) (g^{n_1}(\xi))^{(t_1)} \cdots (g^{n_k}(\xi))^{(t_k)} \equiv a$ ,  $a \in \mathbb{C} \setminus \{0\}$ . We first remark that, from conditions  $a), b)$ , we have that in the case  $n = 0$ , there exists  $i \in \{1, \dots, k\}$  such that  $n_i > t_i$ . Therefore, in both cases ( $n = 0$  and  $n \neq 0$ ), since  $a \neq 0$ , it is easy to see that  $g$  is entire having no zero. So, by Lemma 7,  $g(\xi) = e^{c\xi+d}$ ,  $c \neq 0$ . Then

$$\begin{aligned} g^n(\xi) (g^{n_1}(\xi))^{(t_1)} \cdots (g^{n_k}(\xi))^{(t_k)} &= e^{nc\xi+nd} (e^{n_1c\xi+n_1d})^{(t_1)} \cdots (e^{n_kc\xi+n_kd})^{(t_k)} \\ &= (n_1c)^{t_1} \cdots (n_kc)^{t_k} e^{(n+\sum_{j=1}^k n_j)c\xi+(n+\sum_{j=1}^k n_j)d}. \end{aligned}$$

Then  $(n_1c)^{t_1} \cdots (n_kc)^{t_k} e^{(n+\sum_{j=1}^k n_j)c\xi+(n+\sum_{j=1}^k n_j)d} \equiv a$ , which is impossible. So,  $g^n(\xi) (g^{n_1}(\xi))^{(t_1)} \cdots (g^{n_k}(\xi))^{(t_k)}$  is nonconstant, which proves the claim.

By the assumption of Theorem 1 and by Hurwitz's theorem, for every  $m \in \{1, \dots, q\}$ , all zeros of  $g(\xi)^n (g^{n_1}(\xi))^{(t_1)} \cdots (g^{n_k}(\xi))^{(t_k)} - a_m$  have multiplicity at least  $\ell_m$ .

For any  $j \in \{1, \dots, k\}$ , we have that  $(g^{n_j}(\xi))^{(t_j)}$  is nonconstant. Indeed, if  $(g^{n_j}(\xi))^{(t_j)}$  is constant for some  $j \in \{1, \dots, k\}$ , then since  $n_j \geq t_j$ , and since  $g$  is nonconstant, we get that  $n_j = t_j$  and  $g(\xi) = a\xi + b$ , where  $a, b$  are constants,  $a \neq 0$ . Thus, we can write

$$g(\xi)^n (g^{n_1}(\xi))^{(t_1)} \cdots (g^{n_k}(\xi))^{(t_k)} = c(a\xi + b)^{n+\sum_{j=1}^k (n_j-t_j)},$$

where  $c$  is a nonzero constant. This contradicts to the fact that all zeros of  $g(\xi)^n (g^{n_1}(\xi))^{(t_1)} \cdots (g^{n_k}(\xi))^{(t_k)} - a_m$  have multiplicity at least  $\ell_m \geq 2$  (note that  $a_m \neq 0$ , and that, by condition b) of Theorem 1,  $n + \sum_{j=1}^k (n_j - t_j) > 0$ ). Thus,  $(g^{n_j}(\xi))^{(t_j)}$  is nonconstant, for all  $j \in \{1, \dots, k\}$ .

On the other hand, we can write

$$(g^{n_j})^{(t_j)} = \sum c_{m_0, m_1, \dots, m_{t_j}} g^{m_0} (g')^{m_1} \dots (g^{(t_j)})^{m_{t_j}},$$

$c_{m_0, m_1, \dots, m_{t_j}}$  are constants, and  $m_0, m_1, \dots, m_{t_j}$  are nonnegative integers such that  $m_0 + \dots + m_{t_j} = n_j$ ,  $\sum_{j=1}^{t_j} j m_j = t_j$ . Thus, by an easy computation, we get that  $d(P) = n + \sum_{j=1}^k n_j$ ,  $\theta(P) = \sum_{j=1}^k t_j$ .

Now, we apply Lemma 10 for the differential polynomial

$$P = g(\xi)^n (g^{n_1}(\xi))^{(t_1)} \dots (g^{n_k}(\xi))^{(t_k)}.$$

By Lemma 10, we have (note that, by condition b) of Theorem 1,  $n + \sum_{j=1}^k n_j \geq 2$ )

$$\begin{aligned} T(r, g) &\leq \frac{q \sum_{j=1}^k t_j + 1}{qn + q \sum_{j=1}^k n_j - 1} \bar{N}(r, \frac{1}{g}) \\ &\quad + \frac{1}{qn + q \sum_{j=1}^k n_j - 1} \sum_{m=1}^q \bar{N}(r, \frac{1}{P - a_m}) + o(T(r, g)). \end{aligned} \quad (3.5)$$

For any  $m \in \{1, \dots, q\}$ , we have, by the First Main Theorem,

$$\begin{aligned} \bar{N}(r, \frac{1}{P - a_m}) &= \bar{N}(r, \frac{1}{g^n (g^{n_1})^{(t_1)} \dots (g^{n_k})^{(t_k)} - a_m}) \\ &\leq \frac{1}{\ell_m} N(r, \frac{1}{g^n (g^{n_1})^{(t_1)} \dots (g^{n_k})^{(t_k)} - a_m}) \\ &\leq \frac{1}{\ell_m} T(r, g^n (g^{n_1})^{(t_1)} \dots (g^{n_k})^{(t_k)}) + O(1) \\ &= \frac{1}{\ell_m} m(r, g^n (g^{n_1})^{(t_1)} \dots (g^{n_k})^{(t_k)}) \\ &\quad + \frac{1}{\ell_m} N(r, g^n (g^{n_1})^{(t_1)} \dots (g^{n_k})^{(t_k)}) + O(1). \end{aligned} \quad (3.6)$$

By the Lemma on Logarithmic Derivative and by the First Main Theorem,

$$\begin{aligned}
& m(r, g^n (g^{n_1})^{(t_1)} \dots (g^{n_k})^{(t_k)}) + N(r, g^n (g^{n_1})^{(t_1)} \dots (g^{n_k})^{(t_k)}) \\
& \leq m(r, \frac{g^n (g^{n_1})^{(t_1)} \dots (g^{n_k})^{(t_k)}}{g^n g^{n_1} \dots g^{n_k}}) + m(r, g^n g^{n_1} \dots g^{n_k}) \\
& \quad + N(r, g^n (g^{n_1})^{(t_1)} \dots (g^{n_k})^{(t_k)}) \\
& \leq (n + \sum_{j=1}^k n_j) m(r, g) + N(r, g^n (g^{n_1})^{(t_1)} \dots (g^{n_k})^{(t_k)}) + o(T(r, g)) \\
& = (n + \sum_{j=1}^k n_j) m(r, g) + (n + \sum_{j=1}^k n_j) N(r, g) + (\sum_{j=1}^k t_j) \bar{N}(r, g) + o(T(r, g)) \\
& \leq (n + \sum_{j=1}^k n_j) T(r, g) + (\sum_{j=1}^k t_j) \bar{N}(r, g) + o(T(r, g)). \tag{3.7}
\end{aligned}$$

Combining with (3.6), for all  $m \in \{1, \dots, q\}$  we have

$$\begin{aligned}
\bar{N}(r, \frac{1}{P - a_m}) & \leq \frac{1}{\ell_m} (n + \sum_{j=1}^k n_j) T(r, g) + \frac{1}{\ell_m} (\sum_{j=1}^k t_j) \bar{N}(r, g) + o(T(r, g)) \\
& \leq \frac{1}{\ell_m} (n + \sum_{j=1}^k n_j + \sum_{j=1}^k t_j) T(r, g) + o(T(r, g)). \tag{3.8}
\end{aligned}$$

Therefore, by (3.5) and by the First Main Theorem, we have

$$\begin{aligned}
(qn + q \sum_{j=1}^k n_j - 1) T(r, g) & \leq (q \sum_{j=1}^k t_j + 1) \bar{N}(r, \frac{1}{g}) + \sum_{m=1}^q \bar{N}(r, \frac{1}{P - a_m}) + o(T(r, g)) \\
& \leq (q \sum_{j=1}^k t_j + 1) T(r, g) + (n + \sum_{j=1}^k n_j + \sum_{j=1}^k t_j) (\sum_{m=1}^q \frac{1}{\ell_m}) T(r, g) + o(T(r, g)).
\end{aligned}$$

This implies that

$$\frac{qn + \sum_{j=1}^k q(n_j - t_j) - 2}{n + \sum_{j=1}^k (n_j + t_j)} T(r, g) \leq \sum_{m=1}^q \frac{1}{\ell_m} T(r, g) + o(T(r, g)).$$

Combining with assumption b) we get that  $g$  is constant. This is a contradiction. Hence  $\mathcal{F}$  is a normal family. We have completed the proof of Theorem 1.  $\square$

We can obtain Theorem 3 by an argument similar to the the proof of Theorem 1: We first remark that although condition b) of Theorem 3 is different from condition b) of Theorem 1, whereever it has been used in the proof of Theorem 1 before equation (3.5), the condition b) of Theorem 3 still allows the same conclusion. And from equation (3.5) on we modify as follows : Since  $\mathcal{F}$  is a family of holomorphic functions and by Remark 8,  $g$  is an entire functions. So, similarly to (3.5), by Lemma 10, we have

$$\begin{aligned} T(r, g) &\leq \frac{q \sum_{j=1}^k t_j + 1}{qn + q \sum_{j=1}^k n_j} \overline{N}(r, \frac{1}{g}) + \frac{1}{q(n + \sum_{j=1}^k n_j)} \sum_{m=1}^q \overline{N}(r, \frac{1}{P - a_m}) + o(T(r, g)) \\ &\leq \frac{q \sum_{j=1}^k t_j + 1}{qn + q \sum_{j=1}^k n_j} T(r, g) + \frac{1}{q(n + \sum_{j=1}^k n_j)} \sum_{m=1}^q \overline{N}(r, \frac{1}{P - a_m}) + o(T(r, g)). \end{aligned} \quad (3.9)$$

Since  $g$  is a holomorphic function,  $\overline{N}(r, g) = 0$ . Therefore, by (3.6) and (3.7) (which remain unchanged), we have

$$\overline{N}(r, \frac{1}{P - a_m}) \leq \frac{1}{\ell_m} (n + \sum_{j=1}^k n_j) T(r, g) + o(T(r, g)). \quad (3.10)$$

By (3.9), (3.10), we have

$$\frac{qn + \sum_{j=1}^k q(n_j - t_j) - 1}{n + \sum_{j=1}^k n_j} T(r, g) \leq \sum_{m=1}^q \frac{1}{\ell_m} T(r, g) + o(T(r, g)).$$

Combining with assumption b) of Theorem 3, we get that  $g$  is constant. This is a contradiction. We have completed the proof of Theorem 3.  $\square$

In connection with Remark 5, we note that the proofs of Theorem 1 and Theorem 3 remain valid for the case where the monomial  $f^n(f^{n_1})^{(t_1)} \dots (f^{n_k})^{(t_k)}$  is replaced by the following polynomial

$$f^n(f^{n_1})^{(t_1)} \dots (f^{n_k})^{(t_k)} + \sum_I c_I f^{n_I} (f^{n_{1I}})^{(t_{1I})} \dots (f^{n_{kI}})^{(t_{kI})},$$

where  $c_I$  is a holomorphic function on  $D$ , and  $n_I, n_{jI}, t_{jI}$  are nonnegative integers satisfying

$$\alpha_I := \frac{\sum_{j=1}^k t_{jI}}{n_I + \sum_{j=1}^k n_{jI}} < \alpha := \frac{\sum_{j=1}^k t_j}{n + \sum_{j=1}^k n_j}.$$

In fact, since  $\alpha_I < \alpha$  and by (4.1), we get

$$g_{Iv}(\xi) := \frac{f_v(z_v + \rho_v \xi)}{\rho_v^{\alpha_I}} = \rho_v^{\alpha - \alpha_I} g_v(\xi) \rightarrow 0,$$

spherically uniformly on compact subsets of  $\mathbb{C}$ .

Therefore, similarly to (3.4)

$$\begin{aligned} c_I(z_v + \rho_v \xi) f_v^{n_I}(z_v + \rho_v \xi) (f_v^{n_{1I}})^{(t_{1I})}(z_v + \rho_v \xi) \cdots (f_v^{n_{kI}})^{(t_{kI})}(z_v + \rho_v \xi) \\ = c_I(z_v + \rho_v \xi) g_{Iv}^{n_I}(\xi) (g_{Iv}^{n_{1I}}(\xi))^{(t_{1I})} \cdots (g_{Iv}^{n_{kI}}(\xi))^{(t_{kI})} \rightarrow 0, \end{aligned}$$

spherically uniformly on compact subsets of  $\mathbb{C}$ .

This implies that

$$\begin{aligned} f_v^n(z_v + \rho_v \xi) (f_v^{n_1})^{(t_1)}(z_v + \rho_v \xi) \cdots (f_v^{n_k})^{(t_k)}(z_v + \rho_v \xi) \\ + \sum_I c_I(z_v + \rho_v \xi) f_v^{n_I}(z_v + \rho_v \xi) (f_v^{n_{1I}})^{(t_{1I})}(z_v + \rho_v \xi) \cdots (f_v^{n_{kI}})^{(t_{kI})}(z_v + \rho_v \xi) \\ = g_v^n(\xi) (g_v^{n_1}(\xi))^{(t_1)} \cdots (g_v^{n_k}(\xi))^{(t_k)} \\ + \sum_I c_I(z_v + \rho_v \xi) g_{Iv}^{n_I}(\xi) (g_{Iv}^{n_{1I}}(\xi))^{(t_{1I})} \cdots (g_{Iv}^{n_{kI}}(\xi))^{(t_{kI})} \\ \rightarrow g^n(\xi) (g^{n_1}(\xi))^{(t_1)} \cdots (g^{n_k}(\xi))^{(t_k)}. \end{aligned} \quad (3.11)$$

spherically uniformly on compact subsets of  $\mathbb{C}$ .

We use again the proofs of Theorem 1 and Theorem 3 for the general case above after changing (3.4) by (3.11).  $\square$

## 4 Appendix

Using our methods above, we give a slightly simpler proof of the case of Theorem B above which did not follow from our Corollary 4:

**Theorem 11** ([6], Theorem 3.2, case  $n = k + 1$ ). *Let  $k$  be a positive integer and  $a$  be a nonzero constant. Let  $\mathcal{F}$  be a family of entire functions in a complex domain  $D$  such that for every  $f \in \mathcal{F}$ ,  $(f^{k+1})^{(k)}(z) \neq a$  for all  $z \in D$ . Then  $\mathcal{F}$  is normal on  $D$ .*

In order to prove the above theorem we need the following lemma:

**Lemma 12** ([4]). *Let  $g$  be a transcendental holomorphic function on the complex plane  $\mathbb{C}$ , and  $k$  be a positive integer. Then  $(g^{k+1})^{(k)}$  assumes every nonzero value infinitely often.*

**Proof of Theorem 11.** Without loss the generality, we may assume that  $D$  is the unit disc. Suppose that  $\mathcal{F}$  is not normal at  $z_0 \in D$ . Then, by Lemma 6, for  $\alpha = \frac{k}{k+1}$  there exist

- 1) a real number  $r$ ,  $0 < r < 1$ ,
  - 2) points  $z_v$ ,  $|z_v| < r$ ,  $z_v \rightarrow z_0$ ,
  - 3) positive numbers  $\rho_v$ ,  $\rho_v \rightarrow 0^+$ ,
  - 4) functions  $f_v$ ,  $f_v \in \mathcal{F}$
- such that

$$g_v(\xi) = \frac{f_v(z_v + \rho_v \xi)}{\rho_v^\alpha} \rightarrow g(\xi) \quad (4.1)$$

spherically uniformly on compact subsets of  $\mathbb{C}$ , where  $g(\xi)$  is a non-constant holomorphic function and  $g^\#(\xi) \leq g^\#(0) = 1$ .

Therefore

$$\begin{aligned} (f_v^{k+1})^{(k)}(z_v + \rho_v \xi) &= \left( \frac{f_v(z_v + \rho_v \xi)}{\rho_v^\alpha} \right)^{k+1} \rho_v^{\alpha k} \\ &= (g_v^{k+1}(\xi))^{(k)} \rightarrow (g^{k+1}(\xi))^{(k)} \end{aligned}$$

spherically uniformly on compact subsets of  $\mathbb{C}$ .

By Hurwitz's theorem either  $(g^{k+1})^{(k)} \equiv a$ , either  $(g^{k+1})^{(k)} \neq a$ . On the other hand, it is easy to see that there exists  $z_0$  such that  $(g^{k+1})^{(k)}(z_0) = a$  (the case where  $g$  is a nonconstant polynomial is trivial and the case where  $g$  is transcendental follows from Lemma 12). Hence,  $(g^{k+1})^{(k)} \equiv a$ . Therefore  $g$  has no zero point. Hence, by Lemma 7,  $g(\xi) = e^{c\xi+d}$ ,  $c \neq 0$ . Then  $a \equiv (g^{k+1})^{(k)}(\xi) \equiv ((k+1)c)^k e^{(k+1)(c\xi+d)}$ , which is impossible.  $\square$

## References

- [1] C. T. Chuang, On differential polynomials. In: Analysis of One Complex Variable, World Sci. Publishing, Singapore, 1987, 12-32.
- [2] J. Clunie and W. K. Hayman, *The spherical derivative of integral and meromorphic functions*, Comm. Math. Helv., **40** (1966), 117-148.



- [3] W. K. Hayman, *Meromorphic Functions*, Clarendon Press, Oxford, 1964.
- [4] W. Hennekemper, *Über die Werteverteilung von  $(f^{k+1})^{(k)}$* , Math. Z., **177** (1981), 375-380.
- [5] J. D. Hinchliffe, *On a result of Chuang related to Hayman's Alternative*, Comput. Method. Funct. Theory, **2** (2002), 293-297.
- [6] W. Schwick, *Normal criteria for families of meromorphic functions*, J. Anal. Math., **52** (1989), 241-289.
- [7] X. C. Pang and L. Zalcman, *On theorems of Hayman and Clunie*, New Zealand J. Math., **28** (1999), 71-75.
- [8] L. Zalcman, *Normal families: New perspective*, Bull. Amer. Math. Soc., **35** (1998), 215-230.